arXiv:1207.3616v1 [math.DG] 16 Jul 2012

G_2-STRUCTURES ON EINSTEIN SOLVMANIFOLDS

MARISA FERNÁNDEZ, ANNA FINO AND VICTOR MANERO

Abstract. We study the G_2 analogue of the Goldberg conjecture on non-compact solvmanifolds, i.e. on simply conneted solvable Lie groups no having a lattice. In contrast to the almost-Kähler case we prove that a 7-dimensional solvmanifold cannot admit any left-invariant calibrated G_2-structure φ such that the induced metric g_φ is Einstein, unless g_φ is flat. Moreover, we show that a 7-dimensional (non-flat) Einstein solvmanifold (S, g) of rank one cannot admit any left-invariant cocalibrated G_2-structure φ such that the induced metric $g_\varphi = g$.

1. Introduction

A 7-dimensional smooth manifold M^7 is said to admit a G_2–structure if there is a reduction of the structure group of its frame bundle from $GL(7, \mathbb{R})$ to the exceptional Lie group G_2 which can actually be viewed naturally as a subgroup of $SO(7)$. Therefore a G_2-structure determines a Riemannian metric and an orientation. In fact, one can prove that the presence of a G_2-structure is equivalent to the existence of a certain type of a non-degenerate 3-form φ on the manifold. By [10] a manifold M^7 with a G_2-structure comes equipped with a Riemannian metric g, a cross product P, a 3-form φ, and orientation, which satisfy the relation

$$\varphi(X, Y, Z) = g(P(X, Y), Z).$$

This is exactly analogous to the data of an almost Hermitian manifold, which comes with a Riemannian metric, an almost complex structure J, a 2-form F, and an orientation, which satisfy the relation $F(X, Y) = g(JX, Y)$.

Whenever this 3-form φ is covariantly constant with respect to the LeviCivita connection then the holonomy group is contained in G_2 and the 3-form φ is closed and co-closed.

A G_2-structure is called *calibrated* if the 3-form φ is closed and it can be viewed as the G_2 analogous of an almost-Käher structure in almost Hermitian geometry. By the resuts in [6, 8] no compact 7-dimensional manifold M^7 can support a calibrated G_2-structure φ whose underlying metric g_φ is Einstein unless g_φ has holomomy contained in G_2. This could be considered to be a G_2 analogue of the Goldberg conjecture in almost-Kähler geometry. The result was generalized by R.L.Bryant to calibrated G_2-structures with too tightly pinched Ricci tensor and by R. Cleyton and S. Ivanov to calibrated G_2-structures with divergence-free Weyl tensor.

A non-compact complete Einstein (non-Kähler) almost-Kähler manifold with negative scalar curvature was constructed in [3] and in [13] it was shown that it is an almost-Kähler solvmanifold, that is, a simply connected solvable Lie group S endowed with a left-invariant almost-Kähler structure [13]. In Section 3 we show

---

Date: July 17, 2012.

that in dimension six this is the unique example of Einstein almost-Kähler (non-Kähler) solvmanifold and we classify the 6-dimensional solvmanifolds admitting a left-invariant (non-flat) Kähler-Einstein structure.

A natural problem is then to study the existence of calibrated G_2-structure φ inducing Einstein metrics on non-compact homogeneous Einstein manifolds. All the known examples of non-compact homogeneous Einstein manifolds belong to the class of solvmanifolds, that is, they are simply connected solvable Lie groups S endowed with a left invariant metric (see for instance the survey [18]). A left invariant metric on a Lie group S will be always identified with the inner product $< \cdot, \cdot >$ determined on the Lie algebra $\mathfrak{s}$ of S. According to a long standing conjecture attributed to D. Alekseevskii (see [4, 7.57]), these might exhaust the class of noncompact homogeneous Einstein manifolds.

On the other hand, Lauret in [19] showed that the Einstein solvmanifolds are *standard*, i.e. satisfy the following additional condition: if $\mathfrak{s} = \mathfrak{a} \oplus \mathfrak{n}$ is the orthogonal decomposition of the Lie algebra $\mathfrak{s}$ of S with $\mathfrak{n} = [\mathfrak{s}, \mathfrak{s}]$, then $[\mathfrak{a}, \mathfrak{a}] = 0$.

A left-invariant Ricci-flat metric on solvmanifold is necessarily flat [2], but solvmanifolds can admits incomplete metrics with holonomy G_2 as shown in [11, 7].

In Section 4 by using the classification of 7-dmensional Einstein solvmanifold and obstructions to the existence of calibrated G_2-structures, in contrast to the almost-Kähler case, we prove that a 7-dimensional solvmanifold cannot admit any left-invariant calibrated G_2-structure φ such that the induced metric g_φ is Einstein, unless g_φ is flat.

If φ is co-closed, then the G_2-structure is called *cocalibrated*. In Section 5 we show that a 7-dimensional (non-flat) Einstein solvmanifold (S, g) of rank one cannot admit any left-invariant cocalibrated G_2-structure φ such that the induced metric $g_\varphi = g$.

2. Preliminaries on Einstein solvmanifolds

By [19] all the Einstein solvmanifolds are standard. Standard Einstein solvmanifolds constitute a distinguished class that has been deeply studied by J. Heber, who has derived many remarkable structural and uniqueness results, by assuming only the standard condition (see [12]). In contrast to the compact case, a standard Einstein metric is unique up to isometry and scaling among invariant metrics [12, Theorem E]. The study of standard Einstein solvmanifolds reduces to the rank-one case, that is, to the ones with $\dim \mathfrak{a} = 1$ (see [12, Sections 4.5,4.6]) and everything is determined by the nilpotent Lie algebra $\mathfrak{n} = [\mathfrak{s}, \mathfrak{s}]$. Indeed, a nilpotent Lie algebra $\mathfrak{n}$ is the nilradical of a rank-one Einstein solvmanifold if and only if $\mathfrak{n}$ admits a nilsoliton metric (also called a minimal metric), meaning that its Ricci operator is a multiple of the identity modulo a derivation of $\mathfrak{n}$.

Any standard Einstein solvmanifold is isometric to a solvmanifold whose underlying metric Lie algebra resembles an Iwasawa subalgebra of a semisimple Lie algebra in the sense that ad_A is symmetric and nonzero for any $A \in \mathfrak{a}$, $A \neq 0$. Moreover, if H denotes the mean curvature vector of S (i.e., the only element $H \in \mathfrak{a}$ such that $tr(ad_A) =< A, H >$, for every $A \in \mathfrak{a}$), then the eigenvalues of $ad_H|_{\mathfrak{n}}$ are all positive integers without a common divisor, say $k_1 < \ldots < k_r$. If $d_1, \ldots, d_r$ denote the corresponding multiplicities, then the tuple

$$(k; d) = (k_1 < \ldots < k_r; d_1, \ldots, d_r)$$

is called the *eigenvalue type* of S. It turns out that $\mathbb{R}H \oplus \mathfrak{n}$ is also an Einstein solvmanifold (with just the restriction of $< \cdot, \cdot >$ on it). It is then enough to consider rank-one (i.e. $\dim \mathfrak{a} = 1$) metric solvable Lie algebras as every higher rank Einstein solvmanifold will correspond to a unique rank-one Einstein solvmanifold and certain abelian subalgebra of derivations of $\mathfrak{n}$ containing ad_H. In every dimension, only finitely many eigenvalue types occur.
By [20, Lemma 11], [1]) and [12, Proposition 6.12] it follows that if $(\mathfrak{s}, < \cdot, \cdot >)$ is an Einstein (non-flat) solvable Lie algebra, such that $\dim \mathfrak{a} = m$ and $[\mathfrak{s}, \mathfrak{s}]$ is abelian, then the eigenvalue type is $(1; k)$, with $k = \dim[\mathfrak{s}, \mathfrak{s}] \geq m$.
In the case that $\mathfrak{n}$ is non abelian, it is proved in [17] that any nilpotent Lie algebra of dimension ≤ 5 admits an Einstein solvable extension. In [22] it is shown that the same is true for any of the 34 nilpotent Lie algebras of dimension 6, obtaining then a classification of all 7-dimensional rank-one Einstein solvmanifolds (see Table 2). A classification of 6 and 7-dimensional Einstein solvmanifolds of higher rank can be obtained by [23], where more in general there is a study of Ricci solitons up to dimension 7 on solvmanifolds.

3. Almost-Kähler structures

An almost Hermitian manifold (M, J, g) is called an almost-Kähler manifold if the corresponding Kähler form $F(\cdot, \cdot) = g(\cdot, J\cdot)$ is a closed 2-form In this section we study the existence of Einstein almost-Kähler structures (J, g, F) on 6-dimensional solvmanifolds.
Along all this work, the coefficients appearing in the rank-one Einstein extension of a Lie algebra will be denoted by c_i while the coefficients of the extension up to dimension 6 for almost-Kähler, and up to dimension 7 for G_2 manifolds, will be denoted by f_i.

Theorem 3.1. *A 6-dimensional solvmanifold (S, g) admits a left-invariant Einstein (non-Kähler) almost-Kähler if and only if its Lie algebra $(\mathfrak{s}, g)$ is isometric to the rank two Einstein solvable Lie algebra* (2) *defined below.*
A 6-dimensional solvmanifold (S, g) admits a left-invariant Kähler-Einstein structure if and only if the Lie algebra $(\mathfrak{s}, g)$ is isometric either to the rank one Einstein solvable Lie algebra $\mathfrak{h}_4$ or to the rank two Einstein solvable Lie algebra (1) *or to the rank three Einstein solvable Lie algebra* (3)*; both Lie algebras* (2) *and* (3) *are given below.*

Proof. A 6-dimensional Einstein solvable Lie algebra $(\mathfrak{s}, g)$ is necessarily standard, so one has the orthogonal decomposition (with respect to g)

$$\mathfrak{s} = \mathfrak{n} \oplus \mathfrak{a},$$

with $\mathfrak{n} = [\mathfrak{s}, \mathfrak{s}]$ nilpotent and $\mathfrak{a}$ abelian. We will consider separately the different cases according to the rank if $\mathfrak{s}$, i.e., to the dimension of $\mathfrak{a}$.
If $\dim \mathfrak{a} = 1$ and $\mathfrak{n}$ is abelian, then we know by [12, Proposition 6.12] that $\mathfrak{s}$ has structure equations

$$(ce^{16}, ce^{26}, ce^{36}, ce^{46}, ce^{56}, 0),$$

where c is a non-zero real number. For this Lie algebra we get that any closed 2-form F is degenerate, i.e. satisfies $F^3 = 0$ and so it does not admit symplectic forms.

If $\dim \mathfrak{a} = 1$ but $\mathfrak{n}$ is nilpotent (non-abelian), then $(\mathfrak{s}, g)$ is isometric to one of the solvable Lie algebras $\mathfrak{h}_i$ $(i = 1, \ldots, 8)$ defined below in Table 1, endowed with the Riemannian metric g such that the basis $(e^1, \ldots, e^6)$ is orthonormal.
For $\mathfrak{h}_1$, $\mathfrak{h}_j$, $5 \leq j \leq 8$, we get again that any closed 2-form F is degenerate.

The Lie algebras $\mathfrak{h}_2$ and $\mathfrak{h}_3$ admit symplectic forms. However, one can check that any almost complex structure J on $\mathfrak{h}_i$ $(i = 2, 3)$ is such that $g(\cdot, \cdot) \neq F(\cdot, J\cdot)$.
For $\mathfrak{h}_4$ we get that a symplectic form is

$$F = c_{5,6}e^{12} + c_{1,6}e^{16} + c_{2,6}e^{26} + c_{5,6}e^{34} + c_{3,6}e^{36} + c_{4,6}e^{46} + c_{5,6}e^{56},$$

with $c_{5,6} \neq 0$. The almost complex structures J such that $g(\cdot, \cdot) = F(\cdot, J\cdot)$ are given, with respect to the basis $(e^1, \ldots, e^6)$, by

$$Je_1 = \pm e_2, \quad Je_3 = \pm e_4, \quad Je_5 = \pm e_6$$

and they are integrable. Therefore, (J, g, F) are Kähler-Einsten structures on $\mathfrak{h}_4$.
In order to determine all the 6-dimensional rank-two Einstein solvable Lie algebras, we need first to find the rank-one Einstein solvable extensions $\mathfrak{n}_4 \oplus \mathbb{R} < e_5 >$ of the 4-dimensional nilpotent Lie algebras $\mathfrak{n}_4$. Then we consider the standard solvable Lie algebra $\mathfrak{s}_6 = \mathfrak{n}_4 \oplus \mathfrak{a}$, with $\mathfrak{a} = \mathbb{R} < e_5, e_6 >$ abelian and such that the basis $(e^1, \ldots, e^6)$ is orthonormal. The two cases that we should consider for $\mathfrak{n}_4$ are $(0, 0, e^{12}, 0)$ and $(0, 0, e^{12}, e^{13})$. We find that they have the following rank-one Einstein solvable extensions

$$(\tfrac{1}{2}c_3e^{15}, \tfrac{1}{2}c_3e^{25}, \tfrac{1}{4}\sqrt{22}c_3e^{12} + c_3e^{35}, \tfrac{3}{4}c_3e^{45}, 0),$$

if $\mathfrak{n}_4 = (0, 0, e^{12}, 0)$; and

$$(\tfrac{1}{4}c_4e^{15}, \tfrac{1}{2}c_4e^{25}, \tfrac{1}{2}\sqrt{5}c_4e^{12} + \tfrac{3}{4}c_4e^{35}, \tfrac{1}{2}\sqrt{5}c_4e^{13} + c_4e^{45}, 0),$$

if $\mathfrak{n}_4 = (0, 0, e^{12}, e^{13})$. Now, from the rank-one Einstein solvable extension of $\mathfrak{n}_4 = (0, 0, e^{12}, 0)$, we get the 6-dimensional Einstein solvable Lie algebra of rank two:

$$(1) \qquad \begin{cases} de^1 = 2\sqrt{\frac{2}{33}}Ke^{15} + f_1e^{16} + f_2e^{26}, \\ de^2 = f_2e^{16} + 2\sqrt{\frac{2}{33}}Ke^{25} + f_{10}e^{26}, \\ de^3 = \frac{2}{\sqrt{3}}Ke^{12} + 4\sqrt{\frac{2}{33}}Ke^{35} + (f_1 + f_{10})\,e^{36}, \\ de^4 = \sqrt{\frac{6}{11}}Ke^{45} - 2\,(f_1 + f_{10})\,e^{46}, \\ de^5 = de^6 = 0, \end{cases}$$

where $K = \sqrt{3f_1^2 + 5f_{10}f_1 + f_2^2 + 3f_{10}^2}$, which admits the Kähler-Einstein structures given, in terms of the orthonormal basis $(e^1, \ldots e^6)$, by

$$\begin{aligned} F = {} & \tfrac{1}{2}\sqrt{\tfrac{11}{2}}c_{3,5}e^{12} + c_{1,5}e^{15} + \frac{\sqrt{\frac{33}{2}}e^{16}(f_1c_{1,5}+f_2c_{2,5})}{2B} + c_{2,5}e^{25} + \frac{\sqrt{\frac{33}{2}}e^{26}(f_2c_{1,5}+f_{10}c_{2,5})}{2B} \\ & +c_{3,5}e^{35} + \frac{\sqrt{\frac{33}{2}}(f_1+f_{10})c_{3,5}e^{36}}{4B} + c_{4,5}e^{45} - \frac{\sqrt{\frac{22}{3}}(f_1+f_{10})c_{4,5}e^{46}}{B} + c_{5,6}e^{56}, \end{aligned}$$

$$Je_1 = e_2, Je_2 = -e_1, Je_3 = 2\sqrt{\tfrac{2}{11}}e_5 + \sqrt{\tfrac{3}{11}}e_6, Je_4 = \sqrt{\tfrac{3}{11}}e_5 - 2\sqrt{\tfrac{2}{11}}e_6,$$
$$Je_5 = -2\sqrt{\tfrac{2}{11}}e_3 - \sqrt{\tfrac{3}{11}}e_4, Je_6 = -\sqrt{\tfrac{3}{11}}e_3 + 2\sqrt{\tfrac{2}{11}}e_4,$$

with $(f_1 + f_{10})c_{3,5}^2c_{4,5} \neq 0$.

From the rank-one Einstein solvable extension of $\mathfrak{n}_4 = (0, 0, e^{12}, e^{13})$ we get the 6-dimensional Einstein solvable Lie algebra of rank two:

$$
\begin{cases}
de^1 = \frac{b_4 e^{15}}{4} + \frac{3b_4 e^{16}}{4},\\
de^2 = \frac{b_4 e^{25}}{2} - b_4 e^{26},\\
de^3 = \frac{1}{2}\sqrt{5} b_4 e^{12} + \frac{3b_4 e^{35}}{4} - \frac{b_4 e^{36}}{4},\\
de^4 = \frac{1}{2}\sqrt{5} b_4 e^{13} + b_4 e^{45} + \frac{b_4 e^{46}}{2},\\
de^5 = de^6 = 0,
\end{cases}
\tag{2}
$$

which admit the Einstein (non-Kähler) almost-Kähler structure given by

$$
\begin{aligned}
&F = -2\sqrt{5}c_{3,6}e^{12} + \sqrt{5}c_{4,6}e^{13} + \tfrac{1}{3}c_{1,6}e^{15} + c_{1,6}e^{16} - \tfrac{1}{2}c_{2,6}e^{25} + c_{2,6}e^{26} - 3c_{3,6}e^{35}\\
&\qquad +c_{3,6}e^{36} + 2c_{4,6}e^{45} + c_{4,6}e^{46} + c_{5,6}e^{56},\\
&Je_1 = e_3, Je_3 = -e_1, Je_2 = -\tfrac{1}{\sqrt{5}}e_5 + \tfrac{2}{\sqrt{5}}e_6, Je_4 = \tfrac{2}{\sqrt{5}}e_5 + \tfrac{1}{\sqrt{5}}e_6,\\
&Je_5 = \tfrac{1}{\sqrt{5}}e_2 - \tfrac{2}{\sqrt{5}}e_4, Je_6 = -\tfrac{1}{\sqrt{5}}e_2 - \tfrac{2}{\sqrt{5}}e_4.
\end{aligned}
$$

where $c_{4,6}(4c_{3,6}^2 + c_{2,6}c_{4,6}) \neq 0$. The almost-Kähler is not integrable since

$$N_J(e_1, e_2) = -\sqrt{5}b_4 e_3, N_J(e_1, e_5) = b_4 e_1, N_J(e_1, e_6) = -2b_4 e_1$$

If $\mathfrak{n} = [\mathfrak{s}, \mathfrak{s}]$ is abelian and $\dim \mathfrak{n} = 4$, we have to consider the structure equations

$$
\begin{cases}
de^1 = c_1 e^{15} + f_1 e^{16},\\
de^2 = c_1 e^{25} + f_3 e^{26},\\
de^3 = c_1 e^{35} + f_5 e^{36},\\
de^4 = c_1 e^{45} + f_7 e^{46},\\
de^5 = de^6 = 0
\end{cases}
$$

and then to impose that the metric for which $(e^1, \dots, e^6)$ is orthonormal has to be Einstein and $d^2 e^j = 0$, $j = 1, \dots, 5$. We find that the structure equations are:

$$
\begin{cases}
de^1 = \frac{\sqrt{f_1^2+(f_3+f_4)f_1+f_3^2+f_4^2+f_3f_4}\,e^{15}}{\sqrt{2}} + f_1 e^{16},\\
de^2 = \frac{\sqrt{f_1^2+(f_3+f_4)f_1+f_3^2+f_4^2+f_3f_4}\,e^{25}}{\sqrt{2}} + \left(-f_1 - f_3 - f_4\right) e^{26},\\
de^3 = \frac{\sqrt{f_1^2+(f_3+f_4)f_1+f_3^2+f_4^2+f_3f_4}\,e^{35}}{\sqrt{2}} + f_3 e^{36},\\
de^4 = \frac{\sqrt{f_1^2+(f_3+f_4)f_1+f_3^2+f_4^2+f_3f_4}\,e^{45}}{\sqrt{2}} + f_4 e^{46},\\
de^5 = de^6 = 0.
\end{cases}
$$

and that the corrisponding Lie algebra does not admit any symplectic form.
In order to determine all the 6-dimensional rank-three Einstein solvable Lie algebras, we need first to find the rank-one Einstein solvable extensions $\mathfrak{n}_3 \oplus \mathbb{R} < e_4 >$ of the 3-dimensional nilpotent Lie algebras $\mathfrak{n}_3 = \mathfrak{h}_3$ with structure equations

$$(0, 0, e^{12}).$$

We find the following rank-one Einstein solvable extension

$$(\tfrac{a_3}{2}e^{14}, \tfrac{a_3}{2}e^{24}, a_3 e^{12} + a_3 e^{34}, 0),$$

which is indeed flat.

If $\mathfrak{n} = [\mathfrak{s}, \mathfrak{s}]$ is abelian and $\dim \mathfrak{n} = 3$, we have the Einstein solvable Lie algebra with structure equations

$$(3) \qquad \begin{cases} de^1 = \sqrt{2} f_4 e^{14} - \sqrt{3} f_4 e^{15} + f_4 e^{16}, \\ de^2 = \sqrt{2} f_4 e^{24} + \sqrt{3} f_4 e^{25} + f_4 e^{26}, \\ de^3 = \sqrt{2} f_4 e^{34} - 2 f_4 e^{36}, \\ de^4 = de^5 = de^6 = 0. \end{cases}$$

which admits the almost-Kähler structures given by

$$\begin{aligned} F = {} & \sqrt{2} c_{1,6} e^{14} - \sqrt{3} c_{1,6} e^{15} + c_{1,6} e^{16} + \sqrt{2} c_{2,6} e^{24} + \sqrt{3} c_{2,6} e^{25} + c_{2,6} e^{26} - \tfrac{c_{3,6} e^{34}}{\sqrt{2}} \\ & + c_{3,6} e^{36} + c_{4,5} e^{45} + c_{4,6} e^{46} + c_{5,6} e^{56}, \end{aligned}$$

$$Je_1 = \tfrac{1}{\sqrt{3}} e_4 - \tfrac{1}{\sqrt{2}} e_5 + \tfrac{1}{\sqrt{6}} e_6, Je_2 = \tfrac{1}{\sqrt{3}} e_4 + \tfrac{1}{\sqrt{2}} e_5 + \tfrac{1}{\sqrt{6}} e_6, Je_3 = -\tfrac{1}{\sqrt{3}} e_4 + \sqrt{\tfrac{2}{3}} e_6,$$
$$Je_4 = -\tfrac{1}{\sqrt{3}} e_1 - \tfrac{1}{\sqrt{3}} e_2 + \tfrac{1}{\sqrt{3}} e_3, Je_5 = \tfrac{1}{\sqrt{2}} e_1 - \tfrac{1}{\sqrt{2}} e_2, Je_6 = -\tfrac{1}{\sqrt{6}} e_1 - \tfrac{1}{\sqrt{6}} e_2 - \sqrt{\tfrac{2}{3}} e_3.$$

□

$\mathfrak{s}_6$	6-dimensional Einstein solvable Lie algebras of rank one
$\mathfrak{h}_1$	$(\frac{2}{13} c_4 e^{16}, \frac{2}{13} c_4 e^{26}, \frac{10}{13}\sqrt{3} c_4 e^{12} + \frac{11}{13} c_4 e^{36}, \frac{20}{13} c_4 e^{13} + c_4 e^{46}, \frac{10}{13}\sqrt{3} c_4 e^{14} + \frac{15}{13} c_4 e^{56}, 0)$
$\mathfrak{h}_2$	$(\frac{1}{4} c_4 e^{16}, \frac{1}{4} c_4 e^{26}, \frac{1}{4}\sqrt{30} c_4 e^{12} + \frac{3}{4} c_4 e^{36}, \frac{1}{4}\sqrt{30} c_4 e^{13} + c_4 e^{46}, \frac{1}{2}\sqrt{5} c_4 e^{14} + \frac{1}{2}\sqrt{5} c_4 e^{23} + \frac{5}{4} c_4 e^{56}, 0)$
$\mathfrak{h}_3$	$(\frac{3}{10} c_5 e^{16}, \frac{2}{5} c_5 e^{26}, \frac{3}{5} c_5 e^{36}, \frac{1}{5}\sqrt{30} c_5 e^{12} + \frac{7}{10} c_5 e^{46}, \frac{1}{5}\sqrt{30} c_5 e^{23} + \frac{1}{5}\sqrt{15} c_5 e^{14} + c_5 e^{56}, 0)$
$\mathfrak{h}_4$	$(\frac{1}{2} c_5 e^{16}, \frac{1}{2} c_5 e^{26}, \frac{1}{2} c_5 e^{36}, \frac{1}{2} c_5 e^{46}, c_5 e^{12} + c_5 e^{34} + c_5 e^{56})$
$\mathfrak{h}_5$	$(\frac{1}{2} c_3 e^{16}, \frac{1}{2} c_3 e^{26}, 2 c_3 e^{12} + c_3 e^{36}, \sqrt{3} c_3 e^{13} + \frac{3}{2} c_3 e^{46}, \sqrt{3} c_3 e^{23} + \frac{3}{2} c_3 e^{56}, 0)$
$\mathfrak{h}_6$	$(\frac{1}{3} a_1 e^{16}, \frac{1}{2} a_1 e^{26}, \frac{1}{2} a_1 e^{36}, a_1 e^{12} + \frac{5}{6} a_1 e^{46}, a_1 e^{13} + \frac{5}{6} a_1 e^{56}, 0)$
$\mathfrak{h}_7$	$(\frac{1}{2} c_3 e^{16}, \frac{1}{2} c_3 e^{26}, \frac{1}{2}\sqrt{7} c_3 e^{12} + c_3 e^{36}, \frac{3}{4} c_3 e^{46}, \frac{3}{4} c_3 e^{56}, 0)$
$\mathfrak{h}_8$	$(\frac{1}{4} c_4 e^{16}, \frac{1}{2} c_4 e^{26}, \frac{1}{4}\sqrt{26} c_4 e^{12} + \frac{3}{4} c_4 e^{36}, \frac{1}{4}\sqrt{26} c_4 e^{13} + c_4 e^{46}, \frac{3}{4} c_4 e^{56}, 0)$

Table 1. Rank-one Einstein 6-dimensional solvable Lie algebras

4. Calibrated G_2-structures

In this section we study the existence of calibrated G_2-structures φ on 7-dimensional solvable Lie algebras whose underlying Riemannian metric g_φ is Einstein. We will use the classification of the 7-dimensional Einstein solvable Lie algebras and the following obstructions.

Lemma 4.1. [9] *If there is a non zero vector X in a 7-dimensional Lie algebra $\mathfrak{g}$ such that $(i_X\varphi)^3 = 0$ for all closed 3-form $\varphi \in Z^3(\mathfrak{g}^*)$, then $\mathfrak{g}$ does not admit any calibrated G_2-structure.*

Lemma 4.2. *Let $\mathfrak{g}$ be a 7-dimensional Lie algebra and φ a G_2-structure on $\mathfrak{g}$. Then the bilinear form $g_\varphi : \mathfrak{g} \times \mathfrak{g} \to \mathbb{R}$ defined by*

$$g_\varphi(X, Y) vol = \frac{1}{6}(i_X\varphi \wedge i_Y\varphi \wedge \varphi)$$

has to be a Remannian metric.

Proof. It follows by the fact that in general there is a $1-1$ correspondence between G_2-structures on a 7-manifold and 3-forms φ for which the bilinear form B_φ defined by

$$B_\varphi(X, Y) = (i_X\varphi \wedge i_Y\varphi \wedge \varphi)$$

is positive definite (see [5], [14]). □

Lemma 4.3. *Let* $(\mathfrak{s}, g)$ *be a 7-dimensional Enstein solvable Lie algebra endowed with a* G_2*-structure* φ*, then, for any* $A \in \mathfrak{a} = [\mathfrak{n}, \mathfrak{n}]^\perp$ *such that* $g_\varphi(A, A) = 1$*, the forms*

$$\alpha = i_A\varphi, \quad \beta = \varphi - \alpha \wedge A^*,$$

define an $SU(3)$*-structure on* $(\mathbb{R} < A >)^\perp$*, where by* $A^* \in \mathfrak{s}^*$ *we denote the dual of* A*. So in particular* $\alpha \wedge \beta = 0$ *and* $\alpha^3 \neq 0$*.*

Proof. It follows by Proposition 4.5 in [21]. □

In contrast with the almost-Kähler case, we can prove the following theorem

Theorem 4.4. *A 7-dimensional solvmanifold cannot admit any left-invariant* G_2*-structure* φ *such that* g_φ *is Einstein, unless* g_φ *is flat.*
In particular, if the 7-dimensional Einstein (non-flat) solvmanifold (S, g) *has rank one, then* (S, g) *admits a calibrated* G_2*-structure if and only if the Lie algebra* $\mathfrak{s}$ *of* S *is isometric to the Einstein solvable Lie algebras* $\mathfrak{g}_1$*,* $\mathfrak{g}_4$*,* $\mathfrak{g}_9$*,* $\mathfrak{g}_{18}$*,* $\mathfrak{g}_{28}$ *in Table 2.*

Proof. A 7-dimensional Einstein solvable Lie algebra $(\mathfrak{s}, g)$ is necessarily standard, so one has the orthogonal decomposition (with respect to g)

$$\mathfrak{s} = \mathfrak{n} \oplus \mathfrak{a},$$

with $\mathfrak{n} = [\mathfrak{s}, \mathfrak{s}]$ nilpotent and $\mathfrak{a}$ abelian. We will consider separately the different cases according to the rank if $\mathfrak{s}$, i.e., to the dimension of $\mathfrak{a}$.
If $\dim \mathfrak{a} = 1$ and $\mathfrak{n}$ is abelian, then we know by [12, Proposition 6.12] that $\mathfrak{s}$ has structure equations

$$(ce^{17}, ce^{27}, ce^{37}, ce^{47}, ce^{57}, ce^{67}, 0),$$

where c is a non-zero real number. Computing the generic closed 3-form on $\mathfrak{s}$ it is easy to check that $\mathfrak{s}$ cannot admit any calibrated G_2-structure.
If $\dim \mathfrak{a} = 1$ and $\mathfrak{n}$ is nilpotent (non-abelian), then $(\mathfrak{s}, g)$ is isometric to one of the solvable Lie algebras $\mathfrak{g}_i, i = 1, \dots, 33$, in Table 2, endowed with the Riemannian metric such that the basis $(e^1, \dots, e^7)$ is orthonormal. We may apply Lemma 4.1 with $X = e_6$ to all the Lie algebras $\mathfrak{g}_i, i = 1, \dots, 33$, except to the Lie algebras $\mathfrak{g}_1$, $\mathfrak{g}_4$, $\mathfrak{g}_9$, $\mathfrak{g}_{18}$, $\mathfrak{g}_{28}$, showing in this way that they do not admit any calibrated G_2-structure. For the remaining Lie algebras $\mathfrak{g}_1, \mathfrak{g}_4, \mathfrak{g}_9, \mathfrak{g}_{18}, \mathfrak{g}_{28}$ we first determine the generic closed 3-form φ and then, by applying Lemma 4.3, we impose, that $\alpha \wedge \alpha \neq 0$ and $\alpha \wedge \beta = 0$, where

$$\alpha = i_{e_7}\varphi, \quad \beta = \varphi - e^7 \wedge \beta. \tag{4}$$

Moreover, we have that the closed 3-form φ defines a G_2-structure if and only the matrix associated to the symmetric bilinear form g_φ, with respect to the orthonormal basis $(e^1, \dots, e^7)$, is positive definite. Since the Einstein metric is unique up to scaling, a calibrated G_2-structure induces an Einstein metric if and only if the matrix associated to the symmetric bilinear form g_φ, with respect to the basis $\{e^1, \dots, e^7\}$, is a multiple of the identity matrix. By a direct computation we have that then the Lie algebras $\mathfrak{g}_1, \mathfrak{g}_4, \mathfrak{g}_9, \mathfrak{g}_{18}, \mathfrak{g}_{28}$ admit a calibrated G_2-structure (see Table 3) but they do not admit any calibrated G_2-structure inducing a Einstein (non-flat) metric.
Next, we show that result for the Lie algebra $\mathfrak{g}_{28}$. To this end, we see that any closed 3-form φ on $\mathfrak{g}_{28}$ has the following expression:

$\varphi = c_{1,2}e^{127} - \frac{1}{2}c_{5,6}e^{136} + c_{2,4}e^{137} + \frac{1}{2}c_{5,6}e^{145} - c_{2,3}e^{147} - c_{2,6}e^{157} + c_{2,5}e^{167} + \frac{1}{2}c_{5,6}e^{235}$
$+ c_{2,3}e^{237} + \frac{1}{2}c_{5,6}e^{246} + c_{2,4}e^{247} + c_{2,5}e^{257} + c_{2,6}e^{267} + c_{3,4}e^{347} + c_{3,5}e^{357} + c_{3,6}e^{367}$
$+ c_{3,6}e^{457} - c_{3,5}e^{467} + c_{5,6}e^{567}$,
where $c_{i,j}$ are arbitrary constants.

In this case, one can check that the induced metric by φ can be expressed by the following matrix:

$$G = \begin{pmatrix} -\frac{1}{4}c_{1,2}c_{5,6}^2 & 0 & \frac{1}{4}c_{2,3}c_{5,6}^2 & \frac{1}{4}c_{2,4}c_{5,6}^2 & \frac{1}{4}c_{2,5}c_{5,6}^2 & \frac{1}{4}c_{2,6}c_{5,6}^2 & 0 \\ 0 & -\frac{1}{4}c_{1,2}c_{5,6}^2 & -\frac{1}{4}c_{2,4}c_{5,6}^2 & \frac{1}{4}c_{2,3}c_{5,6}^2 & \frac{1}{4}c_{2,6}c_{5,6}^2 & -\frac{1}{4}c_{2,5}c_{5,6}^2 & 0 \\ \frac{1}{4}c_{2,3}c_{5,6}^2 & -\frac{1}{4}c_{2,4}c_{5,6}^2 & -\frac{1}{4}c_{3,4}c_{5,6}^2 & 0 & \frac{1}{4}c_{3,6}c_{5,6}^2 & -\frac{1}{4}c_{3,5}c_{5,6}^2 & 0 \\ \frac{1}{4}c_{2,4}c_{5,6}^2 & \frac{1}{4}c_{2,3}c_{5,6}^2 & 0 & -\frac{1}{4}c_{3,4}c_{5,6}^2 & -\frac{1}{4}c_{3,5}c_{5,6}^2 & -\frac{1}{4}c_{3,6}c_{5,6}^2 & 0 \\ \frac{1}{4}c_{2,5}c_{5,6}^2 & \frac{1}{4}c_{2,6}c_{5,6}^2 & \frac{1}{4}c_{3,6}c_{5,6}^2 & -\frac{1}{4}c_{3,5}c_{5,6}^2 & \frac{c_{5,6}^3}{4} & 0 & 0 \\ \frac{1}{4}c_{2,6}c_{5,6}^2 & -\frac{1}{4}c_{2,5}c_{5,6}^2 & -\frac{1}{4}c_{3,5}c_{5,6}^2 & -\frac{1}{4}c_{3,6}c_{5,6}^2 & 0 & \frac{c_{5,6}^3}{4} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & g_{7,7} \end{pmatrix},$$

where
$g_{7,7} = -c_{5,6}c_{2,3}^2 + c_{1,2}c_{3,5}^2 + c_{1,2}c_{3,6}^2 + c_{2,5}^2c_{3,4} + c_{2,6}^2c_{3,4} + c_{2,5}\left(2c_{2,3}c_{3,6} - 2c_{2,4}c_{3,5}\right)$
$- 2c_{2,6}\left(c_{2,3}c_{3,5} + c_{2,4}c_{3,6}\right) - c_{2,4}^2c_{5,6} + c_{1,2}c_{3,4}c_{5,6}$. Now we have that the system $G = B \cdot I_7$ does not have solution, for any real number B, where I_7 is the identity matrix. This means that the Lie algebra $\mathfrak{g}_{28}$ does not admit calibrated G_2 structure defining an Einstein metric. However, we can solve 48 from the 49 equations of the system $G = B \cdot I_7$, and we obtain the metric defined by the matrix

$$G = 2\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 4 \end{pmatrix}.$$

Since this matrix is positive definite, the Lie algebra $\mathfrak{g}_{28}$ has a calibrated G_2 form

$$\varphi = -2e^{127} - 2e^{347} - e^{136} + e^{145} + e^{235} + e^{246} + 2e^{567},$$

which induces the metric defined by G.
In order to determine all the 7-dimensional rank two Einstein solvable Lie algebras, we need first to find the rank-one Einstein solvable extensions $\mathfrak{s}_6 = \mathfrak{n}_5 \oplus \mathbb{R} < e_6 >$ of any of the eight 5-dimensional nilpotent Lie algebras $\mathfrak{n}_5$ (see Table 4) and then consider the standard solvable Lie algebra $\mathfrak{s}_7 = \mathfrak{n}_5 \oplus \mathfrak{a}$, with $\mathfrak{a} = \mathbb{R} < e_6, e_7 >$ abelian and such that the basis $(e^1, \ldots, e^7)$ is orthonormal. From $\mathfrak{h}_1$ we get the 7-dimensional Einstein Lie algebra of rank two with structure equations

$$\begin{cases} de^1 = \frac{1}{3}\sqrt{6}ae^{16} + 4ae^{17}, \\ de^2 = \frac{3}{2}\sqrt{6}ae^{26} - 7ae^{27}, \\ de^3 = \frac{5}{3}\sqrt{18}ae^{12} + \frac{11}{6}\sqrt{6}ae^{36} - 3ae^{37}, \\ de^4 = \frac{10}{3}\sqrt{6}ae^{13} + \frac{13}{6}\sqrt{6}ae^{46} + ae^{47}, \\ de^5 = \frac{5}{3}\sqrt{18}ae^{14} + \frac{5}{2}\sqrt{6}ae^{56} + 5ae^{57}, \\ de^6 = de^7 = 0. \end{cases}$$

By computing the generic closed 3-form φ and by using Lemma 4.2 and Lemma 4.3, we get that the matrix associated to g_φ, with respect to the basis $(e^1, \dots, e^7)$, cannot be a multiple of the identity matrix.
From $\mathfrak{h}_2$ we do not get any 7-dimensional Einstein Lie algebra of rank two. From $\mathfrak{h}_3$ we get the 7-dimensional Einstein Lie algebra of rank two with structure equations

$$\begin{cases} de^1 = \frac{1}{7}\sqrt{21}ae^{16} - ae^{17}, \\ de^2 = \frac{4}{21}\sqrt{21}ae^{26} + 2ae^{27}, \\ de^3 = \frac{2}{7}\sqrt{21}ae^{36} - 2ae^{37}, \\ de^4 = \frac{2}{21}\sqrt{30}\sqrt{21}ae^{12} + \frac{1}{3}\sqrt{21}ae^{46} + ae^{47}, \\ de^5 = \frac{2}{21}\sqrt{30}\sqrt{21}ae^{14} + \frac{2}{21}\sqrt{15}\sqrt{21}ae^{23} + \frac{10}{21}\sqrt{21}ae^{56}, \\ de^6 = de^7 = 0. \end{cases}$$

By computing the generic closed 3-form φ and by using Lemma 4.2 and Lemma 4.3, we get that the matrix associated to g_φ, with respect to the basis $(e^1, \dots, e^7)$, cannot be a multiple of the identity matrix. From $\mathfrak{h}_4$ we get the 7-dimensional Einstein Lie algebras of rank two with structure equations

$$\begin{cases} de^1 = \frac{1}{2}\sqrt{f_7^2 + f_2^2 + 2f_3^2 + 2f_4^2 + f_{14}^2 + f_{19}^2}e^{16} - f_7e^{17} + f_2e^{27} + f_3e^{37} + f_4e^{47}, \\ de^2 = \frac{1}{2}\sqrt{f_7^2 + f_2^2 + 2f_3^2 + 2f_4^2 + f_{14}^2 + f_{19}^2}e^{26} + f_2e^{17} + f_7e^{27} + f_4e^{37} - f_3e^{47}, \\ de^3 = \frac{1}{2}\sqrt{f_7^2 + f_2^2 + 2f_3^2 + 2f_4^2 + f_{14}^2 + f_{19}^2}e^{36} + f_3e^{17} + f_4e^{27} - f_{19}e^{37} + f_{14}e^{47}, \\ de^4 = \frac{1}{2}\sqrt{f_7^2 + f_2^2 + 2f_3^2 + 2f_4^2 + f_{14}^2 + f_{19}^2}e^{46} + f_4e^{17} - f_3e^{27} + f_{14}e^{37} + f_{19}e^{47}, \\ de^5 = \sqrt{f_7^2 + f_2^2 + 2f_3^2 + 2f_4^2 + f_{14}^2 + f_{19}^2}(e^{12} + e^{34} + e^{56}), \\ de^6 = de^7 = 0. \end{cases}$$

We may then apply Lemma 4.1 with $X = e_5$.
From $\mathfrak{h}_5$ we get the 7-dimensional Einstein Lie algebras of rank two with structure equations

$$\begin{cases} de^1 = \frac{\sqrt{6}}{6}\sqrt{a^2+b^2}e^{16} + ae^{17} + be^{27}, \\ de^2 = \frac{\sqrt{6}}{6}\sqrt{a^2+b^2}e^{26} + be^{17} - ae^{27}, \\ de^3 = \frac{2\sqrt{6}}{3}\sqrt{a^2+b^2}e^{12} + \frac{\sqrt{6}}{3}\sqrt{a^2+b^2}e^{36}, \\ de^4 = \sqrt{2}\sqrt{a^2+b^2}e^{13} + \frac{\sqrt{6}}{2}\sqrt{a^2+b^2}e^{46} + ae^{47} + be^{57}, \\ de^5 = \sqrt{2}\sqrt{a^2+b^2}e^{23} + \frac{\sqrt{6}}{2}\sqrt{a^2+b^2}e^{56} + be^{47} - ae^{57}, \\ de^6 = de^7 = 0. \end{cases}$$

For these Lie algebras in order to study completely all possibilities we will study separately the two cases $b \neq 0$ and $b = 0$. By computing the generic closed 3-form φ and using Lemma 4.3 we have that the system $g_\varphi(e_i, e_j) - k\delta_i^j = 0$ (where k is a non zero positive real number) with variables the coefficients c_{ijk} of e^{ijk} in φ has no solutions.

From $\mathfrak{h}_6$ we get the two families of 7-dimensional Einstein Lie algebras of rank two with respectively structure equations

$$1)\begin{cases} de^1 = \frac{1}{3}\sqrt{5f_{25}^2+8f_{19}f_{25}+5f_{19}^2+2f_{12}^2}\,e^{16}+2(f_{19}+f_{25})e^{17},\\ de^2 = \frac{1}{2}\sqrt{5f_{25}^2+8f_{19}f_{25}+5f_{19}^2+2f_{12}^2}\,e^{26}-(f_{19}+2f_{25})e^{27}+f_{12}e^{37},\\ de^3 = \frac{1}{2}\sqrt{5f_{25}^2+8f_{19}f_{25}+5f_{19}^2+2f_{12}^2}\,e^{36}+f_{12}e^{27}-(f_{19}+2f_{25})e^{37},\\ de^4 = \sqrt{5f_{25}^2+8f_{19}f_{25}+5f_{19}^2+2f_{12}^2}\,(e^{12}+\frac{5}{6}e^{46})+f_{19}e^{47}+f_{12}e^{57},\\ de^5 = \sqrt{5f_{25}^2+8f_{19}f_{25}+5f_{19}^2+2f_{12}^2}\,(e^{13}+\frac{5}{6}e^{56})+f_{12}e^{47}+f_{25}e^{57},\\ de^6 = de^7 = 0. \end{cases}$$

and

$$2)\begin{cases} de^1 = \sqrt{2}f_{25}e^{16}+4f_{25}e^{17},\\ de^2 = \frac{3}{2}\sqrt{2}f_{25}e^{26}-3f_{25}e^{27}-f_{12}e^{37},\\ de^3 = \frac{3}{2}\sqrt{2}f_{25}e^{36}+f_{12}e^{27}-3f_{25}e^{37},\\ de^4 = 3\sqrt{2}f_{25}e^{12}+\frac{5}{2}\sqrt{2}f_{25}e^{46}+f_{25}e^{47}-f_{12}e^{57},\\ de^5 = 3\sqrt{2}f_{25}e^{13}+\frac{5}{2}\sqrt{2}f_{25}e^{56}+f_{12}e^{47}+f_{25}e^{57},\\ de^6 = de^7 = 0. \end{cases}$$

For 1) we compute first the generic closed 3-forms φ and then, using Lemma 4.3 for $A = e_7$, we impose the condition $\alpha \wedge \beta = 0$. By this condition I get in particular that

$$c_{123}c_{134}(f_{19}+f_{25}) = 0,$$

where by c_{ijk} we denote the coefficient of e^{ijk} in φ. One can immediately exclude the case $c_{134} = 0$, since otherwise the element of the matrix associated to the metric g_φ has to be zero. Then we study separately the cases $c_{123} = 0$ and $f_{19} = -f_{25}$. In both cases we do not find any solution for the system $g_\varphi(e_i, e_j) - k\delta_i^j = 0$. For 2) we study separately the cases $f_{12}f_{25} \neq 0$, $f_{12} = 0$ and $f_{25} = 0$.

In the case $f_{12}f_{25} \neq 0$ we compute first the generic closed 3-forms φ and then, using Lemma 4.3 for $A = e_7$, we impose the condition $\alpha \wedge \alpha \neq 0$, getting the condition $c_{125} \neq 0$. Thus, we take the system $S_{ij} = g_\varphi(e_i, e_j) - k\delta_i^j = 0$ and get the values of $c_{256}, c_{346}, c_{356}$ and c_{236} from $S_{5,5}, S_{3,5}, S_{4,4}$ and $S_{3,4}$. Now $S_{3,3} = -k$, and the system does not admit solution.

In the case $f_{12} = 0$ we first compute the generic closed 3-forms φ and then we use Lemma 4.3 for $A = e_7$, obtaining that $c_{236}, c_{246}, c_{356}$ and c_{456} are all different of zero.

In the case case $f_{25} = 0$ we first compute the generic closed 3-forms φ and then we may apply Lemma 4.1 with $X = e_1, \ldots, e_5$.

From $\mathfrak{h}_7$ we get the four families of 7-dimensional Einstein Lie algebras of rank two with respectively structure equations

$$i)\begin{cases} de^1 = \frac{2}{21}Ke^{16}+(-f_7+f_{13})e^{17}+f_6e^{27},\\ de^2 = \frac{2}{21}Ke^{26}+f_6e^{17}+f_7e^{27},\\ de^3 = \frac{2}{21}K(\sqrt{7}e^{12}+2e^{36})+f_{13}e^{37},\\ de^4 = \frac{1}{7}Ke^{46}-(2f_{13}+f_{25})e^{47}+f_{24}e^{57},\\ de^5 = \frac{1}{7}Ke^{56}+f_{24}e^{47}+f_{25}e^{57},\\ de^6 = de^7 = 0, \end{cases}$$

where $K = \sqrt{21f_7^2 - 21f_7f_{13} + 63f_{13}^2 + 21f_6^2 + 42f_{25}f_{13} + 21f_{25}^2 + 21f_{24}^2}$,

$$ii)\begin{cases} de^1 = \frac{2}{21}\sqrt{21f_7^2 - 21f_7f_{13} + 42f_{13}^2 + 21f_6^2}e^{16} + (-f_7 + f_{13})e^{17} + f_6e^{27}, \\ de^2 = \frac{2}{21}\sqrt{21f_7^2 - 21f_7f_{13} + 42f_{13}^2 + 21f_6^2}e^{26} + f_6e^{17} + f_7e^{27}, \\ de^3 = \frac{2}{21}\sqrt{21f_7^2 - 21f_7f_{13} + 42f_{13}^2 + 21f_6^2}(\sqrt{7}e^{12} + 2e^{36}) + f_{13}e^{37}, \\ de^4 = \frac{1}{7}\sqrt{21f_7^2 - 21f_7f_{13} + 42f_{13}^2 + 21f_6^2}e^{46} - f_{13}e^{47} - f_{24}e^{57}, \\ de^5 = \frac{1}{7}\sqrt{21f_7^2 - 21f_7f_{13} + 42f_{13}^2 + 21f_6^2}e^{56} + f_{24}e^{47} - f_{13}e^{57}, \\ de^6 = de^7 = 0. \end{cases}$$

$$iii)\begin{cases} de^1 = \frac{1}{21}\sqrt{231f_{13}^2 + 168f_{13}f_{25} + 84f_{25}^2 + 84f_{24}^2}e^{16} + \frac{1}{2}f_{13}e^{17} - f_6e^{27}, \\ de^2 = \frac{1}{21}\sqrt{231f_{13}^2 + 168f_{13}f_{25} + 84f_{25}^2 + 84f_{24}^2}e^{26} + f_6e^{17} + \frac{1}{2}f_{13}e^{27}, \\ \frac{1}{21}\sqrt{231f_{13}^2 + 168f_{13}f_{25} + 84f_{25}^2 + 84f_{24}^2}(\sqrt{7}e^{12} + 2e^{36}) + f_{13}e^{37}, \\ de^4 = \frac{1}{14}\sqrt{231f_{13}^2 + 168f_{13}f_{25} + 84f_{25}^2 + 84f_{24}^2}e^{46} - (2f_{13} + f_{25})e^{47} + f_{24}e^{57}, \\ de^5 = \frac{1}{14}\sqrt{231f_{13}^2 + 168f_{13}f_{25} + 84f_{25}^2 + 84f_{24}^2}e^{56} + f_{24}e^{47} + f_{25}e^{57}, \\ de^6 = de^7 = 0. \end{cases}$$

$$iv)\begin{cases} de^1 = \frac{1}{3}\sqrt{3}f_{25}e^{16} - \frac{1}{2}f_{25}e^{17} - f_6e^{27}, \\ de^2 = \frac{1}{3}\sqrt{3}f_{25}e^{26} + f_6e^{17} - \frac{1}{2}f_{25}e^{27}, \\ de^3 = \frac{1}{3}\sqrt{3}f_{25}(\sqrt{7}e^{12} + 2e^{36}) - f_{25}e^{37}, \\ de^4 = \frac{1}{2}\sqrt{3}f_{25}e^{46} + f_{25}e^{47} - f_{24}e^{57}, \\ de^5 = \frac{1}{2}\sqrt{3}f_{25}e^{46} + f_{24}e^{47} + f_{25}e^{57}, \\ de^6 = de^7 = 0. \end{cases}$$

For all of them after computing the generic closed 3-forms we may apply Lemma 4.1 with $X = e_3$.

From $\mathfrak{h}_8$ we get the 7-dimensional Einstein Lie algebras of rank two with structure equations

$$\begin{cases} de^1 = \frac{1}{39}\sqrt{390f_{13}^2 - 78f_{13}f_{19} + 156f_{19}^2}e^{16} + (-f_{13} + f_{19})e^{17}, \\ de^2 = \frac{1}{39}\sqrt{390f_{13}^2 - 78f_{13}f_{19} + 156f_{19}^2}e^{26} + (2f_{13} - f_{19})e^{27}, \\ de^3 = \frac{1}{39}\sqrt{390f_{13}^2 - 78f_{13}f_{19} + 156f_{19}^2}(\sqrt{26}e^{12} + 3e^{36}) + f_{13}e^{37}, \\ de^4 = \frac{1}{39}\sqrt{390f_{13}^2 - 78f_{13}f_{19} + 156f_{19}^2}(\sqrt{26}e^{13} + 4e^{46}) + f_{19}e^{47}, \\ de^5 = \frac{1}{13}\sqrt{390f_{13}^2 - 78f_{13}f_{19} + 156f_{19}^2}e^{56} - (2f_{13} + f_{19})e^{57}, \\ de^6 = de^7 = 0. \end{cases}$$

If $[\mathfrak{s}, \mathfrak{s}] = \mathfrak{n}$ is abelian and $\dim \mathfrak{n} = 5$, we have to consider the structure equations

$$\begin{cases} de^1 = c_1e^{16} + f_1e^{17}, \\ de^1 = c_1e^{26} + f_7e^{27}, \\ de^3 = c_1e^{36} + f_{13}e^{37}, \\ de^4 = c_1e^{46} + f_{19}e^{47} \\ de^5 = c_1e^{56} + f_{25}e^{57}, \\ de^6 = de^7 = 0. \end{cases}$$

By imposing that $\mathfrak{s}$ is a Einstein Lie algebra (the metric is the one for which $(e^1, \ldots, e^7)$ is orthonormal), we get the Lie algebra with structure equations

$$\begin{cases} de^1 = Ke^{16} + (-f_7 - f_{13} - f_{19})e^{17}, \\ de^1 = Ke^{26} + f_7 e^{27}, \\ de^3 = Ke^{36} + f_{13} e^{37}, \\ de^4 = Ke^{46} + f_{19} e^{47} \\ de^5 = Ke^{56}, \\ de^6 = de^7 = 0, \end{cases}$$

where $K = \sqrt{10f_7^2 + 10f_7 f_{13} + 10f_7 f_{19} + 10f_{13}^2 + 10f_{13}f_{19} + 10f_{19}^2}$.
We study separately the cases $f_{13}f_{19} \neq 0$, $f_{13} = 0$ and $f_{19} = 0$.
In the case $f_{13}f_{19} \neq 0$ using Lemma 4.3 (i.e. $\alpha \wedge \alpha \neq 0$, with $A = e_7$) we may suppose $c_{124}c_{135} \neq 0$ for the generic closed 3-form. Now, we consider the system $S_{ij} = g_\varphi(e_i, e_j) - k\delta_i^j = 0$. We take $c_{1,2,5}, c_{1,2,3}$ and k from $S_{4,5} = 0 = S_{3,4}$ and $S_{2,2} = 0$. In the new system we can conclude that from equations $S_{2,4} = 0$ and $S_{3,5} =$ that there is no solution. Indeed from the two equations follows that $f_{13} = -\frac{19}{5} f_{19}$ and $f_{13} = \frac{7}{31} f_{19}$, which is a contradiction since $f_{13}f_{19} \neq 0$.
In the case $f_{13} = 0$ using Lemma 4.3 we may suppose $c_{1,3,5}c_{3,4,7} \neq 0$ for the generic closed 3-form. Then we get $c_{2,5,7}, A$ and $c_{2,3,7}$ from $S_{5,5} = 0 = S_{5,3} = S_{2,3}$. The new system satisfies:

$$S_{4,4} = \frac{7c_{135}\left(49c_{1,2,5}^2 + 152c_{3,4,7}^2\right)}{76\sqrt{78}}.$$

In the case $f_{19} = 0$ using Lemma 4.3 we may suppose $c_{1,3,5}c_{3,4,7} \neq 0$ for the generic closed 3-form.. Then we get $c_{1,2,3}, c_{2,5,6}, c_{2,3,7}$ and $c_{1,3,5}$ from $S_{3,4} = 0 = S_{2,3} = S_{2,4}$ and $S_{3,3} = 0$. The new system satisfies:

$$S_{4,4} = -\frac{959322c_{1,2,5}^2 c_{3,4,7}^4 + 59711k^2}{59711k}.$$

If $[\mathfrak{s}, \mathfrak{s}] = \mathfrak{n}$ is abelian and $\dim \mathfrak{n} = 5$, we have to consider the structure equations

$$\begin{cases} de^1 = c_1 e^{16} + f_1 e^{17}, \\ de^1 = c_1 e^{26} + f_7 e^{27}, \\ de^3 = c_1 e^{36} + f_{13} e^{37}, \\ de^4 = c_1 e^{46} + f_{19} e^{47} \\ de^5 = c_1 e^{56} + f_{25} e^{57}, \\ de^6 = de^7 = 0. \end{cases}$$

By imposing that $\mathfrak{s}$ is a Einstein Lie algebra (the metric is the one for which $(e^1, \dots, e^7)$ is orthonormal), we get the Lie algebras with structure equations

$$\begin{cases} de^1 = Be^{16} + (-f_7 - f_{13} - f_{19})e^{17}, \\ de^1 = Be^{26} + f_7 e^{27}, \\ de^3 = Be^{36} + f_{13} e^{37}, \\ de^4 = Be^{46} + f_{19} e^{47} \\ de^5 = Be^{56}, \\ de^6 = de^7 = 0, \end{cases}$$

where $B = \sqrt{10f_7^2 + 10f_7 f_{13} + 10f_7 f_{19} + 10f_{13}^2 + 10f_{13}f_{19} + 10f_{19}^2}$. For these Lie algebras we first compute the generic closed 3-forms φ and then we may apply Lemma 4.1 with $X = e_1, \dots, e_5$.

In order to determine all the 7-dimensional rank three Einstein solvable Lie algebras, we need first to find the rank-one Einstein solvable extensions $\mathfrak{n}_4 \oplus \mathbb{R} < e_5 >$ of the two 5-dimensional nilpotent Lie algebras $\mathfrak{n}_4$ and then consider the standard solvable Lie algebra $\mathfrak{s}_6 = \mathfrak{n}_4 \oplus \mathfrak{a}$, with $\mathfrak{a} = \mathbb{R} < e_5, e_6, e_7 >$ abelian and such that the basis $(e^1, \dots, e^7)$ is orthonormal. For any of the nilpotent Lie algebras $\mathfrak{n}_4$ we find the following rank-one Einstein solvable extensions

$$\begin{array}{ll} 1) & (\frac{1}{4}ce^{15}, \frac{1}{2}ce^{25}, \frac{1}{2}\sqrt{5}ce^{12} + \frac{3}{4}ce^{35}, \frac{1}{2}\sqrt{5}ce^{13} + ce^{45}, 0) \\ 2) & (\frac{1}{2}ce^{15}, \frac{1}{2}ce^{25}, \frac{1}{4}\sqrt{22}ce^{12} + ce^{35}, \frac{3}{4}ce^{45}, 0) \end{array}$$

From 1) we do not get any 7-dimensional Einstein Lie algebra of rank three. From 2) From 1) we do not get any 7-dimensional Einstein Lie algebra of rank three. From 2) we get the 7-dimensional Einstein Lie algebra of rank three with structure equations

$$\begin{cases} de^1 = \frac{1}{2}c_3e^{15} - (f_{10} + f_{28})e^{16} + f_2e^{26} + (-f_{14} + f_{23})e^{17} + f_6e^{27}, \\ de^2 = \frac{1}{2}c_3e^{25} + f_9e^{16} + f_{10}e^{26} + f_{13}e^{17} + f_{14}e^{27}, \\ de^3 = \frac{1}{4}\sqrt{22}c_3e^{12} + c_3e^{35} - \frac{1}{2}f_{28}e^{36} + f_{23}e^{37}, \\ de^4 = \frac{3}{4}c_3e^{45} + f_{28}e^{46} - 2f_{23}e^{47}, \\ de^5 = de^6 = de^7 = 0. \end{cases}$$

satisfying ones of the following conditions:

(i) $c_3 = \sqrt{\frac{32f_{14}^2 - 32f_{14}f_{23} + 96f_{23}^2 + 32f_{13}^2}{33}}$, $f_2 = f_9 = \pm\sqrt{\frac{11}{4f_{14}^2 - 4f_{14}f_{23} + f_{23}^2 + 4f_{13}^2}} f_{13}f_{23}$, $f_6 = f_{13}, f_{10} = \mp\frac{1}{11}\sqrt{\frac{11}{4f_{14}^2 - 4f_{14}f_{23} + f_{23}^2 + 4f_{13}^2}}(-13f_{14}f_{23} + 6f_{23}^2 + 2f_{14}^2 + 2f_{13}^2)$, $f_{28} = \pm\frac{2}{11}\sqrt{\frac{11}{4f_{14}^2 - 4f_{14}f_{23} + f_{23}^2 + 4f_{13}^2}}(4f_{14}^2 - 4f_{14}f_{23} + f_{23}^2 + 4f_{13}^2)$

(ii) $c_3 = 2\sqrt{\frac{2}{3}}f_{23}, f_2 = \pm\frac{1}{2}\sqrt{11f_{23}^2 - 4f_{10}^2}, f_6 = f_{13} = f_{28} = 0, f_{14} = \frac{1}{2}f_{23}$,

For the case (i) we consider the generic closed 3-form φ and by using all the time that $c_3 \neq 0$ and the condition $\alpha \wedge \beta = 0$ (with $\alpha = i_{e_7}\varphi$). By imposing the the vanishing of $g_\varphi(e_3, e_4)$ and the condition $g_\varphi(e_3, e_3) \neq 0$ we have always that either $g_\varphi(e_2, e_2) = 0$ or $g_\varphi(e_3, e_3) = 0$ so we cannot have calibrated G_2-structures associated to the Einstein metric.
For the case (ii) we start only to impose the conditions

$$c_3 = 2\sqrt{\frac{2}{3}}f_{23}, f_6 = f_{13} = f_{28} = 0, f_{14} = \frac{1}{2}f_{23},$$

one needs for the Einstein condition still to impose that $f_2 = \pm\frac{1}{2}\sqrt{11f_{23}^2 - 4f_{10}^2}$. We consider the generic closed 3-form φ by using all the time that $f_{23} \neq 0$ (since $c_3 \neq 0$) and we impose $\alpha \wedge \beta = 0$, where $\alpha = i_{e_7}\varphi$. By using that $g_\varphi(e_3, e_3) \neq 0$ and the equations

$$g_\varphi(e_3, e_4) = g_\varphi(e_2, e_3) = g_\varphi(e_1, e_3) = 0.$$

Studying separately the solutions of the above system we show that no calibrated G_2-structure can induce the Einstein metric.

If $\mathfrak{n} = [\mathfrak{s}, \mathfrak{s}]$ is abelian and $\dim \mathfrak{n} = 4$, we get the Einstein solvable Lie algebras of rank three with the structure equations

$$\begin{cases} de^1 = c_1 e^{15} + f_1 e^{16} + f_2 e^{17}, \\ de^1 = c_1 e^{25} + f_3 e^{26} + f_4 e^{27}, \\ de^3 = c_1 e^{35} + f_5 e^{36} + f_6 e^{37}, \\ de^4 = c_1 e^{45} + f_7 e^{46} + f_8 e^{47}, \\ de^5 = de^6 = de^7 = 0 \end{cases}$$

satisfying the conditions

$$\begin{array}{l} f_7 = -f_1 - f_3 - f_5, \quad f_8 = -f_2 - f_4 - f_6, \\ 2f_1f_2 + 2f_3f_4 + 2f_5f_6 + f_2f_3 + f_2f_5 + f_1f_4 + f_4f_5 + f_1f_6 + f_3f_6 = 0, \\ 2f_1^2 + 2f_3^2 + 2f_5^2 + 2f_1f_3 + 2f_1f_5 + 2f_3f_5 = 4c_1^2. \end{array}$$

We impose for the generic 3-form φ the conditions $d\varphi = 0$ and $\alpha \wedge \beta = 0$ (with $\alpha = i_{e_7}\varphi$), using all the time that $c_1 \neq 0$. Then we consider the equations

$$g_\varphi(e_1, e_2) = g_\varphi(e_1, e_3) = g_\varphi(e_1, e_4) = g_\varphi(e_2, e_3) = 0 \tag{5}$$

By the conditions

$$g_\varphi(e_1, e_1) \neq 0, \quad g_\varphi(e_2, e_2) \neq 0, \quad g_\varphi(e_3, e_3) \neq 0, \quad g_\varphi(e_4, e_4) \neq 0$$

we get respectively

$$c_{125}c_{135}c_{145} \neq 0, \quad c_{125}c_{235}c_{245} \neq 0, \quad c_{135}c_{235}c_{345} \neq 0, \quad c_{145}c_{245}c_{345} \neq 0.$$

In all the solutions of (5) we have always that either $c_{125} = 0$ or $c_{135} = 0$ or $c_{145} = 0$, which is not possible. □

5. Cocalibrated G_2-structures

In this section we study the existence of cocalibrated G_2-structures φ on 7-dimensional rank-one Einstein solvable Lie algebras $(\mathfrak{s}, g)$ whose underlying Riemannian metric $g_\varphi = g$.

We will use the classification of the 7-dimensional rank one Einstein solvable Lie algebras (see Table 2), Lemma 4.2 and 4.3, together with the following obstruction.

Lemma 5.1. *Let $(\mathfrak{g}, g)$ be a 7-dimensional metric Lie algebra. If for every closed 4-form $\Psi \in Z^4(\mathfrak{g}^*)$ there exists $X \in \mathfrak{g}$ such that $(i_X(*\Psi))^3 = 0$, then $\mathfrak{g}$ does not admit a cocalibrated G_2 structure inducing the metric g.*

Proof. It is sufficient to prove that if a 3-form φ defines a G_2 structure on $\mathfrak{g}$ then for any $X \in \mathfrak{g}$ we have

$$\iota_X(*\varphi) \wedge \varphi \neq 0,$$

where $*$ is the Hodge star operator with respect to the metric g_φ associated to φ. Since the 3-form φ defines a G_2 structure on $\mathfrak{g}$,then there exist a basis of $\mathfrak{g}$ $\{f_1, \cdots, f_7\}$ (which is orthonormal with respect to g_φ) such that

$$\varphi = f^{124} + f^{235} + f^{346} + f^{457} + f^{156} + f^{267} + f^{137},$$

where $(f^1, \cdots, f^7)$ is the basis of $\mathfrak{g}^*$ dual to $(f_1, \cdots, f_7)$. Taking the Hodge $*$ operator with respect to $(f_1, \cdots, f_7)$ we have

$$*\varphi = -f^{3567} + f^{1467} - f^{1257} + f^{1236} + f^{2347} - f^{1345} - f^{2456}.$$

Thus, writing ι_X the contraction by X,

$$\iota_{f_1}(*\varphi) = f^{467} - f^{257} + f^{236} - f^{345}$$

and

$$\iota_{f_1}(*\varphi) \wedge \varphi = 4f^{234567}.$$

Similarly, we have

$$\begin{aligned}
&\iota_{f_2}(*\varphi) = f^{157} - f^{136} + f^{347} - f^{456}, \quad \iota_{f_2}(*\varphi) \wedge \varphi = -4f^{134567};\\
&\iota_{f_3}(*\varphi) = -f^{567} + f^{126} - f^{247} + f^{145}, \quad \iota_{f_3}(*\varphi) \wedge \varphi = 4f^{124567};\\
&\iota_{f_4}(*\varphi) = -f^{167} + f^{237} - f^{135} + f^{256}, \quad \iota_{f_4}(*\varphi) \wedge \varphi = -4f^{123567};\\
&\iota_{f_5}(*\varphi) = f^{367} - f^{127} + f^{134} - f^{246}, \quad \iota_{f_5}(*\varphi) \wedge \varphi = 4f^{123467};\\
&\iota_{f_6}(*\varphi) = -f^{357} + f^{147} - f^{123} + f^{245}, \quad \iota_{f_6}(*\varphi) \wedge \varphi = -4f^{123457};\\
&\iota_{f_7}(*\varphi) = f^{356} - f^{146} + f^{125} - f^{234}, \quad \iota_{f_7}(*\varphi) \wedge \varphi = 4f^{123456}.
\end{aligned}$$

In general, for $i = 1, \cdots 7$, we see that

$$\iota_{f_i}(*\varphi) \wedge \varphi = (-1)^{i+1} 4 f^{12\cdots(i-1)(i+1)\cdots 7},$$

which is a non-zero 6-form for any $i = 1, \ldots, 7$. □

Lemma 5.2. *Let $(\mathfrak{g}, g)$ be a 7-dimensional metric Lie algebra. If for any φ coclosed 3-form on $\mathfrak{g}$, the differential form $\tau_3 = *d\varphi|_{\Lambda^3_{27}\mathfrak{g}^*}$ satisfies the conditions*

$$\varphi \wedge \tau_3 \neq 0 \ \textit{or}\ (*\varphi) \wedge \tau_3 \neq 0$$

then $\mathfrak{g}$ does not admit cocalibrated G_2 structure inducing the metric g.

Proof. The expression of the differential and the codifferential of a G_2 form φ are given in terms of the intrinsic torsion forms by

$$\begin{aligned}
d\varphi &= \tau_0 * \varphi + 3\tau_1 \wedge \varphi + *\tau_3\\
d * \varphi &= 4\tau_1 \wedge *\varphi + \tau_2 \wedge \varphi.
\end{aligned}$$

with $\tau_0 \in \Lambda^0\mathfrak{g}^*$, $\tau_1 \in \Lambda^1\mathfrak{g}^*$, $\tau_2 \in \Lambda^2_{14}\mathfrak{g}^*$ and $\tau_3 \in \Lambda^3_{27}\mathfrak{g}^*$. The cocalibrated condition $d * \varphi = 0$ implies

$$d\varphi = \tau_0 * \varphi + *\tau_3.$$

Since

$$\begin{aligned}
\Lambda^3_{27}\mathfrak{g}^* &= \{\rho \in \Lambda^3\mathfrak{g}^* | \rho \wedge \varphi = 0 = \rho \wedge *\varphi\},\\
\Lambda^4_{27}\mathfrak{g}^* &= \{\gamma \in \Lambda^4\mathfrak{g}^* | \gamma \wedge \varphi = 0 = \gamma \wedge *\varphi\}
\end{aligned}$$

it follows that

$$d\varphi \wedge \varphi = \tau_0 |\varphi|^2 e^{1234567}.$$

Therefore

$$\tau_3 = - * (d\varphi - \tau_0 * \varphi).$$

Now as $\tau_3 \in \Lambda^3_{27}\mathfrak{g}^*$ the conditions

$$\tau_3 \wedge \varphi = 0, \quad \tau_3 \wedge *\varphi = 0$$

must be fulfilled. □

We know already that a 7-dimensional Enstein solvable Lie algebra cannot admit nearly-parallel G_2-structures since the scalar curvature has to be positive. For the co-calibrated G_2-structures we can prove the following

Theorem 5.3. *A 7-dimensional (nonflat) rank-one Einstein solvmanifold (S, g) cannot admit any left-invariant cocalibrated G_2-structure φ such that $g_\varphi = g$.*

Proof. For a 7-dimensional rank-one Einstein solvable Lie algebra $(\mathfrak{s}, g)$ we have the orthogonal decomposition (with respect to the Einstein metric g)

$$\mathfrak{s} = \mathfrak{n}_6 \oplus \mathfrak{a},$$

with $\mathfrak{n}_6 = [\mathfrak{s}, \mathfrak{s}]$ a 6-dimensional nilpotent Lie algebra and $\mathfrak{a} = \mathbb{R} < e_7 >$ abelian. If $\mathfrak{n}$ is abelian, then we know by [12, Proposition 6.12] that $\mathfrak{s}$ has structure equations

$$(ce^{17}, ce^{27}, ce^{37}, ce^{47}, ce^{57}, ce^{67}, 0),$$

where c is a non-zero real number. Computing the generic co-closed 4-form on $\mathfrak{s}$ it easy to check that $\mathfrak{s}$ cannot admit any co-calibrated G_2-structure g_φ such that $g_\varphi = g$.

If $\mathfrak{n}_6$ is nilpotent (non-abelian), then $(\mathfrak{s}, g)$ is isometric to one of the solvable Lie algebras $\mathfrak{g}_i, i = 1, \dots, 33$, in Table 2, endowed with the Riemannian metric such that the basis $(e^1, \dots, e^7)$ is orthonormal. We may apply Lemma 5.1 with $X = e_7$ to the Lie algebras $\mathfrak{g}_3$, $\mathfrak{g}_{13}$, $\mathfrak{g}_{23}$ and $\mathfrak{g}_j$, $25 \leq j \leq 33$, showing in this way that they do not admit any cocalibrated G_2-structure φ such that $g_\varphi = g$. For the Lie algebras:

$$\mathfrak{g}_1, \mathfrak{g}_2, \mathfrak{g}_4, \mathfrak{g}_5, \mathfrak{g}_6, \mathfrak{g}_{20}$$

we first determine the generic co-closed 3-form φ and then, we compute the values of τ_0, of the 3-form τ_3. We have that $\tau_3 \wedge \varphi \neq 0$ unless $\varphi = 0$ and so by applying Lemma 5.2 we have that the Lie algebras do not admit a cocalibrated G_2 structure inducing an Einstein metric. For the Lie algebras

$$\mathfrak{g}_7, \mathfrak{g}_8, \mathfrak{g}_9, \mathfrak{g}_{10}, \mathfrak{g}_{12}, \mathfrak{g}_{14}, \mathfrak{g}_{15}, \mathfrak{g}_{16}, \mathfrak{g}_{17}, \mathfrak{g}_{18}, \mathfrak{g}_{19}, \mathfrak{g}_{21}, \mathfrak{g}_{22}, \mathfrak{g}_{24} \tag{6}$$

we first determine the generic co-closed 3-form φ and then, by applying Lemma 4.3, we impose, that $\alpha \wedge \alpha \neq 0$ and $\alpha \wedge \beta = 0$, where α and β are given in (4). Moreover, we have that the closed 3-form φ defines a G_2-structure if and only the matrix associated to the symmetric bilinear form g_φ, with respect to the orthonormal basis $(e^1, \dots, e^7)$, is positive definite. Since the Einstein metric is unique up to scaling, a calibrated G_2-structure induces an Einstein metric if and only if the matrix associated to the symmetric bilinear form g_φ, with respect to the basis $\{e^1, \dots, e^7\}$, is a multiple of the identity matrix. By a direct computation we have thus that the Lie algebras (6) cannot admit any co-calibrated G_2-structures inducing an Einstein metric. □

$\mathfrak{g}_1$	$\left(\frac{a}{2}e^{17}, ae^{27}, \sqrt{13}ae^{12} + \frac{3}{2}ae^{37}, 4ae^{13} + 2ae^{47}, 2\sqrt{3}e^{14} + 2ae^{23} + \frac{5}{2}ae^{57}, -\sqrt{13}ae^{25} + 2\sqrt{3}ae^{34} + \frac{7}{2}ae^{67}, 0\right)$
$\mathfrak{g}_2$	$\left(-\frac{\sqrt{21}}{42}ae^{17}, -\frac{\sqrt{21}}{14}ae^{27}, ae^{12} - 2\frac{\sqrt{21}}{21}ae^{37}, -2\frac{\sqrt{3}}{3}ae^{13} - 5\frac{\sqrt{21}}{42}ae^{47}, ae^{14} - \frac{\sqrt{21}}{7}ae^{57}, ae^{34} - ae^{25} - 3\frac{\sqrt{21}}{14}ae^{67}, 0\right)$
$\mathfrak{g}_3$	$\left(-\frac{\sqrt{14}}{56}ae^{17}, -9\frac{\sqrt{14}}{56}ae^{27}, ae^{12} - 5\frac{\sqrt{14}}{28}ae^{37}, \frac{\sqrt{6}}{2}ae^{13} - 11\frac{\sqrt{14}}{56}ae^{47}, \frac{\sqrt{6}}{2}ae^{14} - 3\frac{\sqrt{14}}{14}ae^{57}, ae^{15} - 13\frac{\sqrt{14}}{56}ae^{67}, 0\right)$
$\mathfrak{g}_4$	$\left(ae^{17}, 2ae^{27}, 2\sqrt{7}e^{12} + 3ae^{37}, \frac{6\sqrt{154}}{11}ae^{13} + 4ae^{47}, 2\sqrt{7}ae^{14} + 2\frac{\sqrt{1155}}{11}ae^{23} + 5ae^{57}, 2\frac{\sqrt{1155}}{11}ae^{15} + 5\frac{\sqrt{154}}{11}ae^{24} + 6ae^{67}, 0\right)$
$\mathfrak{g}_5$	$\left(ae^{17}, 3ae^{27}, 2\sqrt{14}ae^{12} + 4ae^{37}, 2\sqrt{15}ae^{13} + 5ae^{47}, 6\sqrt{2}ae^{14} + 6ae^{57}, 4\sqrt{2}ae^{15} + 2\sqrt{15}ae^{23} + 7ae^{67}, 0\right)$
$\mathfrak{g}_6$	$\left(\frac{a}{2}e^{17}, ae^{27}, \sqrt{10}ae^{12} + \frac{3}{2}ae^{37}, \sqrt{10}ae^{13} + 2ae^{47}, \sqrt{10}ae^{23} + \frac{5}{2}ae^{57}, \sqrt{10}ae^{14} + \frac{5}{2}ae^{67}, 0\right)$
$\mathfrak{g}_7$	$\left(ae^{17}, ae^{27}, 4ae^{12} + 2ae^{37}, 2\sqrt{5}ae^{13} + 3ae^{47}, 2\sqrt{5}ae^{23} + 3ae^{57}, 4ae^{14} - 4ae^{24}a + 4e^{67}, 0\right)$
$\mathfrak{g}_8$	$\left(ae^{17}, ae^{27}, 4ae^{12} + 2ae^{37}, 2\sqrt{5}ae^{13} + 3ae^{47}, 2\sqrt{5}ae^{23} + 3ae^{57}, 4ae^{14} + 4ae^{24}a + 4e^{67}, 0\right)$
$\mathfrak{g}_9$	$\left(\frac{-3}{14}ae^{17}, \frac{-11}{28}ae^{27}, \frac{-3}{7}ae^{37}, \frac{\sqrt{5}}{2}ae^{12} + \frac{-17}{28}ae^{47}, ae^{14} - ae^{23} + \frac{-23}{28}ae^{57}, ae^{34} + \frac{\sqrt{5}}{2}ae^{15} - \frac{29}{28}ae^{67}, 0\right)$
$\mathfrak{g}_{10}$	$\left(\frac{4}{9}ae^{17}, ae^{27}, \frac{4}{3}ae^{37}, \frac{2\sqrt{114}}{9}ae^{12} + \frac{13}{9}ae^{47}, \frac{2}{9}\sqrt{190}ae^{14} + \frac{17}{9}ae^{57}, \frac{2\sqrt{114}}{9}ae^{15} + \frac{2\sqrt{114}}{9}ae^{23}a + \frac{7}{3}e^{67}, 0\right)$
$\mathfrak{g}_{11}$	$\left(\frac{a}{3}e^{17}, \frac{2}{3}ae^{27}, \frac{10\sqrt{7}}{21}ae^{12} + ae^{37}, \frac{4\sqrt{42}}{21}ae^{12} + ae^{47}, \frac{4\sqrt{105}}{21}ae^{13} + \frac{2\sqrt{70}}{21}ae^{14} + \frac{4}{3}ae^{57}, \frac{2\sqrt{6}}{3}ae^{15} + \frac{2\sqrt{7}}{3}ae^{24}a + \frac{5}{3}e^{67}, 0\right)$
$\mathfrak{g}_{12}$	$\left(\frac{a}{2}e^{17}, ae^{27}, \frac{11}{6}ae^{37}, \frac{2\sqrt{21}}{3}ae^{12} + \frac{3}{2}ae^{47}, \frac{2\sqrt{21}}{3}ae^{14} + 2ae^{57}, \frac{2\sqrt{14}}{3}ae^{15} + \frac{2\sqrt{14}}{3}ae^{24}a + \frac{5}{2}e^{67}, 0\right)$
$\mathfrak{g}_{13}$	$\left(\frac{a}{9}e^{17}, ae^{27}, \frac{4}{3}ae^{37}, \frac{2\sqrt{93}}{9}ae^{12} + \frac{33}{27}ae^{47}, \frac{4\sqrt{31}}{9}ae^{14} + \frac{39}{27}ae^{57}, \frac{2\sqrt{93}}{9}ae^{15} + \frac{5}{3}e^{67}, 0\right)$
$\mathfrak{g}_{14}$	$\left(\frac{a}{2}e^{17}, ae^{27}, \frac{3}{4}ae^{37}, \frac{\sqrt{21}}{2}ae^{12} + \frac{3}{2}ae^{47}, \frac{\sqrt{14}}{2}ae^{13} + \frac{5}{4}ae^{57}, \frac{\sqrt{14}}{2}ae^{14} + \frac{\sqrt{21}}{2}ae^{35} + 2ae^{67}, 0\right)$
$\mathfrak{g}_{15}$	$\left(ae^{17}, ae^{27}, ae^{37}, \sqrt{10}ae^{12} + 2ae^{47}, \sqrt{10}ae^{23} + 2ae^{57}, \sqrt{10}ae^{14} + \sqrt{10}ae^{35} + 3ae^{67}, 0\right)$
$\mathfrak{g}_{16}$	$\left(ae^{17}, ae^{27}, ae^{37}, \sqrt{10}ae^{12} + 2ae^{47}, \sqrt{10}ae^{23} + 2ae^{57}, \sqrt{10}ae^{14} - \sqrt{10}ae^{35} + 3ae^{67}, 0\right)$
$\mathfrak{g}_{17}$	$\left(ae^{17}, ae^{27}, \frac{12}{5}ae^{37}, \frac{4}{5}\sqrt{31}ae^{12} + 2ae^{47}, \frac{2}{5}\sqrt{93}ae^{14} + 3ae^{57}, \frac{2}{5}\sqrt{93}ae^{24} + 3ae^{67}, 0\right)$
$\mathfrak{g}_{18}$	$\left(ae^{17}, ae^{27}, 2ae^{37}, 4ae^{12} + 2ae^{47}, 2ae^{13} - 2\sqrt{3}ae^{24} + 3ae^{57}, 2\sqrt{3}ae^{14} + 2ae^{23} + 3ae^{67}, 0\right)$
$\mathfrak{g}_{19}$	$\left(5ae^{17}, 6ae^{27}, 12ae^{37}, 2\sqrt{134}ae^{12} + 11ae^{47}, \sqrt{402}ae^{14} + 16ae^{57}, \sqrt{134}ae^{13} - \sqrt{402}ae^{24} + 17ae^{67}, 0\right)$
$\mathfrak{g}_{20}$	$\left(ae^{17}, ae^{27}, 2ae^{12} + 2ae^{37}, 2\sqrt{3}ae^{12} + 2ae^{47}, 4ae^{14} + 3ae^{57}, 2ae^{24} + 2\sqrt{3}ae^{23} + 3ae^{67}, 0\right)$
$\mathfrak{g}_{21}$	$\left(3ae^{17}, 5ae^{27}, 6ae^{37}, 2\sqrt{42}ae^{12} + 8ae^{47}, 2\sqrt{21}ae^{13} + 9ae^{57}, 2\sqrt{42}ae^{14} + 2\sqrt{21}ae^{23} + 11ae^{67}, 0\right)$
$\mathfrak{g}_{22}$	$\left(6ae^{17}, 5ae^{27}, 9ae^{37}, 2\sqrt{93}ae^{12} + 11ae^{47}, 2\sqrt{93}ae^{13} + 15ae^{57}, 4\sqrt{31}ae^{24} + 16ae^{67}, 0\right)$
$\mathfrak{g}_{23}$	$\left(ae^{17}, \frac{5}{2}ae^{27}, 3ae^{37}, \sqrt{37}ae^{12} + \frac{7}{2}ae^{47}, \frac{\sqrt{74}}{2}ae^{13} + 4ae^{57}, \sqrt{37}ae^{14} + \frac{9}{2}ae^{67}, 0\right)$
$\mathfrak{g}_{24}$	$\left(ae^{17}, ae^{27}, ae^{37}, \sqrt{6}ae^{12} + 2ae^{47}, \sqrt{6}ae^{13} + 2ae^{57}, \sqrt{6}ae^{23} + 2ae^{67}, 0\right)$
$\mathfrak{g}_{25}$	$\left(\frac{5\sqrt{31}}{124}ae^{17}, \frac{2\sqrt{31}}{31}ae^{27}, \frac{9\sqrt{31}}{124}ae^{37}, \frac{9\sqrt{31}}{124}ae^{47}, ae^{12} + \frac{13\sqrt{31}}{124}ae^{57}, -\frac{\sqrt{3}}{2}ae^{34} - \frac{\sqrt{3}}{2}ae^{15} + \frac{9\sqrt{31}}{62}ae^{67}, 0\right)$
$\mathfrak{g}_{26}$	$\left(ae^{17}, 2ae^{27}, 3ae^{37}, 3ae^{47}, 4\sqrt{2}ae^{12} + 3ae^{57}, 4\sqrt{2}ae^{15} + 4ae^{67}, 0\right)$
$\mathfrak{g}_{27}$	$\left(ae^{17}, \frac{3}{4}ae^{27}, \frac{7}{4}ae^{37}, \frac{3}{2}ae^{47}, \frac{\sqrt{148}}{4}ae^{12} + \frac{7}{4}ae^{57}, \frac{\sqrt{74}}{4}ae^{14} + \frac{\sqrt{37}}{2}ae^{25} + \frac{5}{2}ae^{67}, 0\right)$
$\mathfrak{g}_{28}$	$\left(ae^{17}, ae^{27}, ae^{37}, ae^{47}, 2ae^{13} - 2ae^{24} + 2ae^{57}, 2ae^{14} + 2ae^{23} + 2ae^{67}, 0\right)$
$\mathfrak{g}_{29}$	$\left(ae^{17}, ae^{27}, \frac{4}{3}ae^{37}, \frac{4}{3}ae^{47}, \sqrt{6}ae^{12} + 2ae^{57}, \sqrt{6}ae^{14} + \sqrt{6}ae^{23} + \sqrt{7}3ae^{67}, 0\right)$
$\mathfrak{g}_{30}$	$\left(\frac{a}{2}e^{17}, \frac{a}{2}e^{27}, \frac{a}{2}e^{37}, \frac{a}{2}e^{47}, \sqrt{2}ae^{12} + ae^{57}, \sqrt{2}ae^{34} + ae^{67}, 0\right)$
$\mathfrak{g}_{31}$	$\left(\frac{\sqrt{11}}{11}ae^{17}, \frac{3\sqrt{11}}{22}ae^{27}, \frac{3\sqrt{11}}{22}ae^{37}, \frac{2\sqrt{11}}{11}ae^{47}, \frac{5\sqrt{11}}{22}aae^{57}, ae^{13} + \frac{5\sqrt{11}}{22}ae^{67}, 0\right)$
$\mathfrak{g}_{32}$	$\left(\frac{a}{2}e^{17}, \frac{a}{2}e^{27}, \frac{a}{2}e^{37}, \frac{a}{2}e^{47}, \frac{2}{3}ae^{57}, \frac{\sqrt{11}}{3}ae^{12} + \frac{\sqrt{11}}{3}ae^{34} + ae^{67}, 0\right)$
$\mathfrak{g}_{33}$	$\left(\frac{a}{2}e^{17}, \frac{a}{2}e^{27}, \frac{3}{4}ae^{37}, \frac{3}{4}ae^{47}, \frac{3}{4}ae^{57}, \frac{\sqrt{34}}{4+}ae^{12} + +ae^{67}, 0\right)$

Table 2. Rank-one Einstein 7-dimensional solvable Lie algebras

$\mathfrak{s}_7$	Calibrated G_2-structure
$\mathfrak{g}_1$	$\varphi = \frac{1}{432}\left(1440 - 128\sqrt{3}\right)e^{123} + \frac{\sqrt{13}e^{125}}{2} - \frac{\left(13312\sqrt{3}-748800\right)e^{127}}{44928\sqrt{3}}$ $+\frac{8e^{135}}{9} - 2e^{137} - \frac{e^{146}}{\sqrt{3}} - \sqrt{3}e^{147} + 10e^{157} - e^{167} - \frac{e^{236}}{3} + e^{237}$ $+\frac{1}{576}\left(1440 + 128\sqrt{3}\right)e^{247} + \frac{e^{267}}{\sqrt{3}} + \frac{e^{345}}{\sqrt{3}} - e^{357} - e^{457} + e^{567}$
$\mathfrak{g}_4$	$\varphi = -\frac{7e^{125}}{2\sqrt{5}} + e^{137} - \frac{7e^{146}}{13} - e^{147} + \frac{e^{167}}{2} + \frac{7e^{236}}{13} - e^{237}$ $+2e^{247} - e^{267} + \frac{7e^{345}}{13} + \frac{e^{357}}{2} - e^{457} - e^{567}$
$\mathfrak{g}_9$	$\varphi = -\frac{7e^{125}}{2\sqrt{5}} + e^{137} - \frac{7e^{146}}{13} - e^{147} + \frac{e^{167}}{2} + \frac{7e^{236}}{13}$ $-e^{237} + 2e^{247} - e^{267} + \frac{7e^{345}}{13} + \frac{e^{357}}{2} - e^{457} - e^{567}$
$\mathfrak{g}_{18}$	$\varphi = e^{123} - e^{127} - e^{136} + \sqrt{3}e^{145}$ $+3e^{167} + e^{235} + \sqrt{3}e^{246} - \frac{e^{347}}{2} + 3e^{567}$
$\mathfrak{g}_{28}$	$\varphi = -2e^{127} - 2e^{347} - e^{136} + e^{145} + e^{235} + e^{246} + 2e^{567}$

Table 3. Calibrated G_2-structures on rank-one Einstein solvable Lie algebras.

Acknowledgments. This work has been partially supported through Project MICINN (Spain) MTM2011-28326-C02-02, Project UPV/EHU ref. UFI11/52 and by GNSAGA of INdAM.

References

[1] Alekseevskii D.V. *Homogeneous Riemannian spaces of negative curvature*, Math. USSR. Sb., **25** (1975), 87–109.

[2] Alekseevskii D.V., Kimelfeld B.N. *Structure of homogeneous Riemannian spaces with zero Ricci curvature*, Functional Anal. Appl., **9** (1975), 97–102.

[3] Apostolov V. , Draghici T., Moroianu A. *A splitting theorem for Kähler manifolds whose Ricci tensor have constant eigenvalues*, Internat. J. Math. **12** (2001) 769–789.

[4] Besse A. *Einstein manifolds*, 1987, Springer, Berlin, Heidelberg, New York.

[5] Bryant R. L. *Metrics with exceptional holonomy*, Ann. of Math. (2) **126** (1987), no. 3 525–576.

[6] Bryant R. *Some remarks on G_2-structures*, in Proceedings of Gökova Geometry-Topology Conference 2005 75-109 Gkova Geometry/Topology Conference (GGT), Gökova (2006).

[7] Chiossi S. , Fino A. *Conformally parallel G_2 structures on a class of solvmanifolds*, Math. Z. **252** (2006), 825–848.

[8] Cleyton R., Ivanov S. *On the geometry of closed G_2-structures*, Commun. Math. Phys. **270** (2007), 53–67.

[9] Conti D., Fernandez M. *Nilmanifolds with a calibrated G_2-structure*, Differential Geom. Appl. **29** (2011), no. 4, 493–506.

[10] Fernández M., Gray A. *Riemannian manifolds with structure group G_2*, Ann. Mat. Pura Appl. (4) **132** (1982) 19–45.

[11] Gibbons, G.W., L, H., Pope, C.N., Stelle, K.S. *Supersymmetric domain walls frommetrics of special holonomy*, Nuclear Phys. B **623** (2002), 3–46.

[12] Heber J. *Noncompact homogeneous Einstein spaces*, Invent. Math., **133** (1998), 279–352.

[13] Hirobe K. Oguro T., Sekigawa K., *A remark on an example of a* 6*-dimensional Einstein almost-Kähler manifold*, J. Geom. **88** (2008) 70–74.

[14] Hitchin N. J. *The geometry of three-forms in six and seven dimensions*, math/0010054.

[15] Lauret J. *Ricci soliton homogeneous nilmanifolds*, Math. Annalen, **319** (2001), 715–733.

[16] Lauret J. *Standard Einstein solvmanifolds as critical points*, Quart. J. Math. **52** (2001), 463–470.

[17] Lauret J. *Finding Einstein solvmanifolds by a variational method*, Math. Z. **241** (2002), 83–99.
[18] Lauret J. *Einstein solvmanifolds and nilsolitons*, in New Developments in Lie Theory and Geometry, Contemp. Math. 491, Amer. Math. Soc., Providence, RI, 2009, pp. 1–35.
[19] Lauret J. *Einstein solvmanifolds are standard*, Annals Math. **172** (2010), 1859–1877.
[20] Y. G. Nikonorov, *Noncompact Homogeneous Einstein* 5*-Manifolds*, Geometriae Dedicata (2005) **113**, 107–143.
[21] Schulte-Hengesbach F. *Half-flat structures on Lie groups*, PhD Thesis (2010), Hamburg, available at http://www.math.uni-hamburg.de/home/schulte-hengesbach/diss.pdf.
[22] Will C. *Rank-one Einstein solvmanifolds of dimension* 7, Diff. Geom. Appl. **19** (2003), 307–318.
[23] Will C. *The space of solvsolitons in low dimensions*, Ann. Global Anal. Geom. **40** (2011), 291–309.

Universidad del País Vasco, Facultad de Ciencia y Tecnología, Departamento de Matemáticas, Apartado 644, 48080 Bilbao, Spain.
`marisa.fernandez@ehu.es`
`victormanuel.manero@ehu.es`

Dipartimento di Matematica, Università di Torino, Via Carlo Alberto 10, Torino, Italy.
`annamaria.fino@unito.it`